

Cardiovascular support
Arterial / Cholesterol support
CoQ10
400 mg



Description

Coenzyme Q10, also known as ubiquinone, is present in almost all cells and is essential for mitochondrial energy production.¹⁻³ The highest concentration of Coenzyme Q10 (CoQ10) is found in the heart muscle, where constant chemical energy availability is imperative.¹⁻⁴ In addition to supporting cardiac function through energy production, CoQ10 also serves as a fat-soluble antioxidant, providing vascular structures and other tissues with protection against damages from free radicals and oxidative stress.^{1-4,9} Vitamin E has been added to the formula for additional, synergistic antioxidant protection.

Features & Benefits

- Provides Antioxidant Protection (lipophilic)
- Assists with Cellular Energy Production
- Promotes Cardiovascular Health
- Supports Healthy Heart Muscle Function
- May promote Healthy Blood Pressure already within the normal range
- Supports Healthy Muscular Function
- Replenishes Healthy Blood Levels of CoQ10

Suggested Usage

As a dietary supplement, take one softgel 1-3 times daily with a meal or as directed by your healthcare practitioner.

Allergen Checklist

Contains no sugar, salt, starch, yeast, wheat, gluten, soy, milk, egg, shellfish or preservatives.

Cautions / Interactions

See page 2

Technical Summary

Coenzyme Q10 is a naturally occurring fat-soluble vitamin-like quinone, also called ubiquinone as it is “ubiquitous” or present in all eukaryotic cells. CoQ10 is a mitochondrial coenzyme essential for the production of ATP.^{1-3,8,29,33,39,41} Being at the core of cellular energy processes, it assumes importance in tissues with high energy requirements, such as cardiac cells which seem to be sensitive to low CoQ10 production.^{14,27,28,29} In the past decades, CoQ10 has been a popular research subject, especially concerning its antioxidative properties protecting against peroxidation of lipid membranes. By improving antioxidant status and cellular bioenergetics, CoQ10 supplementation may play a role in promoting heart health.^{4,29,30,31,32,33} Indeed, mounting evidence derived from experimental as well as clinical studies suggests that supplemental CoQ10 may possess cardiosupportive,^{1-13,31-34} cytoprotective,^{1-3,6-8,15} and neuroprotective^{1-3,6-8,15,34} properties. Emerging scientific works also indicate that supplemental CoQ10 may promote healthy aging and longevity.³⁵ In the human body, the fat-soluble antioxidant tocopherol (vitamin E) often works in concert with CoQ10.³⁶

Mechanisms of Action

Clinical studies have demonstrated that supplemental CoQ10 is able to raise blood levels in healthy individuals,³⁷ as well as in those affected by potential deficiency states, such as patients taking certain medications or aging individuals.^{5-7,9,11-13,24,31-34,42,43,46,48-53} Experts consider CoQ10's benzoquinone ring its functional group, which is able to assume three alternate redox states: the fully oxidized or ubiquinone (Q) form; the univalently reduced ($1e^- + 1H^+$) ubisemiquinone ($\bullet QH$), a semi-radical; and the fully reduced ($2e^- + 2H^+$) ubiquinol form. Experts hypothesize that the polyisoprenyl chain facilitates the stability of the molecule within the hydrophobic lipid bilayer. In addition, the length of the CoQ isoprenoid chain may affect the mobility, intermolecular interaction with membrane proteins, and auto-oxidizability.³⁸ The physiological roles of CoQ10 in biological systems are most well characterized in the inner mitochondrial membrane, where its main functions are: (i) carrier of electrons from respiratory complexes I and II to complex III, (ii) generation of superoxide anion radical by auto-oxidation of ubisemiquinone and (iii) quenching of free radicals.³⁹ Thus, CoQ10 plays a key role in mitochondrial bioenergetics, while also functioning as a potent antioxidant.²⁹ It is the only lipid soluble antioxidant synthesized endogenously.

Continued on page 2

Clinical Applications

Based on the currently available research, those desiring to promote cardiovascular health, support the body's energy generating processes, antioxidant defense mechanisms, healthy aging and well-being would benefit from taking CoQ10 supplements. Emerging science also suggests that CoQ10 supplements might be of value for athletes,^{15,45} syndromes associated with sudden decline in auditory function,^{46,47,48} as well as patients with different diagnoses undergoing surgery.^{49,50,51,52} Patients regularly taking therapeutic prescription drugs or undergoing surgery are encouraged to discuss with their physician if CoQ10-supplements might be beneficial for them. Likewise, physicians caring for patients requiring certain prescription drugs are encouraged to review potential drug-nutrient interactions to assess the value of adding CoQ10 supplements.




FIGURE 1 Coenzyme Q Redox Stages: fully reduced ubiquinol form (CoQH₂), the radical semiquinone intermediate (CoQH•), and the fully oxidized ubiquinone form (CoQ).

<http://ead.univ-angers.fr/~jaspard/Page2/COURS/Zsuite/1Respiration/Z999suite/3ChaineRespiratoire/3Figures/5CxesChaineRespir/4Ubiquinone.gif>

Complementary Products

Consider taking this product in combination with Protocol for Life Balance® **Omega-3 (P1656)**, **Ortho-E™ (P0950)**, **Alpha-Lipoic Acid (P3042)**, **Nattokinase (P3140)**.

Supplement Facts

Serving Size 1 Softgel

P3198

Amount Per Serving	
Vitamin E (as d-alpha Tocopherol)	30 IU
Coenzyme Q₁₀ (CoQ₁₀)	400 mg
Soy Lecithin	35 mg

Other Ingredients: Rice Bran Oil, Softgel Capsule (gelatin, glycerin, water, carob) and Beeswax.

Contains no: sugar, salt, starch, yeast, wheat, gluten, corn, milk, egg shellfish or preservatives

Mechanisms of Action Continued

Its reduced form, CoQH₂, inhibits protein and DNA oxidation, as well as lipid peroxidation of cell membrane structures and lipoproteins present in the circulation. Clinical studies have revealed that supplementation with CoQ10 results in increased levels of CoQH₂ within circulating lipoproteins and increased resistance of human low-density lipoproteins (LDL) to the initiation of lipid peroxidation.^{40,41} Experimental data also suggests that CoQ10 may affect the activity of endothelium bound extra cellular Superoxide Dismutase (ecSOD), which is considered a biomarker of vascular function.^{9,29} Indeed, recent clinical studies have demonstrated that CoQ10 administration can support healthy endothelial function.^{42,43} Furthermore, recent scientific data revealed that CoQ10 may affect the expression of genes involved in human cell signaling, metabolism, and transport.⁴⁴

Cautions / Interactions

Coenzyme Q10 supplements may interact with certain medications, such as anticoagulants,^{16,17} and selected chemotherapeutic agents.¹⁸⁻²⁰ Preliminary evidence suggests that doxorubicin's (Adriamycin) cardiotoxicity may be attributed to its interference with CoQ10-synthesis in the heart.¹⁸ Theoretically, taking CoQ10 in supplement form may be of benefit for those taking this medication. However, there is concern that antioxidants might influence some medical treatments, such as chemotherapy (e.g., cyclophosphamide, Cytoxan) and radiation therapy.^{19,20} Furthermore, due to CoQ10's influence on blood pressure,^{5,11-13} it may have additive effects when used concurrently with antihypertensive drugs; accordingly, caution is advised with its combined use. In recent years, statin drugs have been identified to lower CoQ10 levels in the body;²¹⁻²³ current research suggests that CoQ10 supplementation may be useful when taking these medications.^{24,25,26} Consult with your healthcare professional if you are taking any of these agents.

References on page 3

Formulated by doctors and clinical scientists exclusively for licensed healthcare practitioners. Manufactured in an A-rated Good Manufacturing (GMP) Certified facility.

PROTOCOL
FOR LIFE BALANCE™

Protocol For Life Balance™

Manufactured by NHG, Bloomingdale, IL 60108 Made in the U.S.A.

Toll Free 877.PROTO10 / Fax 800.886.1045 / www.protocolforlife.com

Healthy Patients, Satisfied Physicians™



REFERENCES

1. Bonakdar RA, Guarneri E. Coenzyme Q10. *Am Fam Physician*. 2005;15;72(6):1065-70. Review.
2. Chow CK. Dietary coenzyme Q10 and mitochondrial status. *Methods Enzymol*. 2004;382:105-12.
3. Nohl N, Staniek K, Kozlov AV, Gille L. The biomolecule ubiquinone exerts a variety of biological functions. *BioFactors*. 2003 18:23-31.
4. Jones K, Hughes K, Mischley L, McKenna DJ. Coenzyme Q-10 and cardiovascular health. *Altern Ther Health Med*. 2004;10(1):22-30.
5. Sander S, Coleman CI, Patel AA, et al. The impact of coenzyme Q10 on systolic function in patients with chronic heart failure. *J Card Fail*. 2006;12(6):464-72.
6. Makhija N, Sendasgupta C, Kiran U, et al. Influence of CoQ10 on autonomic nervous activity and energy metabolism during exercise in healthy subjects. *J Cardiothorac Vasc Anesth*. 2008;22(6):832-9.
7. Zheng A, Moritani T. Influence of CoQ10 on autonomic nervous activity and energy metabolism during exercise in healthy subjects. *J Nutr Sci Vitaminol* 2008;54(4):286-90.
8. Mizuno K, Tanaka M, Nozaki S, et al. Antifatigue effects of coenzyme Q10 during physical fatigue. *Nutrition*. 2008;24(4):293-9.
9. Tiano L, Belardinelli R, Carnevali P, et al. Effect of coenzyme Q10 administration on endothelial function and extracellular superoxide dismutase in patients with ischaemic heart disease: a double-blind, randomized controlled study. *Eur Heart J* 2007;28:2249-55.
10. Belardinelli R, Muçaj A, Lacialprice F, et al. Coenzyme Q10 and exercise training in chronic heart failure. *Eur Heart J*. 2006;27(22):2675-81.
11. Burke BE, Neuenschwander R, Olson RD. Randomized, double-blind, placebo-controlled trial of coenzyme Q10 in isolated systolic hypertension. *South Med J* 2001;94:1112-7.
12. Hodgson JM, Watts GF, Playford DA, et al. Coenzyme Q10 improves blood pressure and glycaemic control: a controlled trial in subjects with type 2 diabetes. *Eur J Clin Nutr* 2002;56:1137-42.
13. Langsjoen P, Willis R, Folkers K. Treatment of essential hypertension with coenzyme Q10. *Mol Aspects Med* 1994;S265-72.
14. Sacconi S, Trevisson E, Salvati L, et al. Coenzyme Q(10) is frequently reduced in muscle of patients with mitochondrial myopathy. *Neuromuscul Disord*. 2010;20(1):44-8.
15. Kon M, Tanabe K, Akimoto T, et al. Reducing exercise-induced muscular injury in kendo athletes with supplementation of coenzyme Q10. *Br J Nutr*. 2008;100(4):903-9.
16. Spigset O. Reduced effect of warfarin caused by ubidecarenone. *Lancet* 1994;334:1372-3.
17. Heck AM, DeWitt BA, Lukes AL. Potential interactions between alternative therapies and warfarin. *Am J Health Syst Pharm* 2000;57:1221-7.
18. Eaton S, Skinner R, Hale JP, et al. Plasma coenzyme Q(10) in children and adolescents undergoing doxorubicin therapy. *Clin Chim Acta* 2000;302:1-9.
19. Portakal O, Ozkaya O, Erden Inal M, et al. Coenzyme Q10 concentrations and antioxidant status in tissues of breast cancer patients. *Clin Biochem* 2000;33:279-84.
20. Lund EL, Quistorff B, Spang-Thomsen M, Kristjansen PE. Effect of radiation therapy on small-cell lung cancer is reduced by ubiquinone intake. *Folia Microbiol* 1998;43:505-6.
21. Folkers K, Langsjoen P, Willis R, et al. Lovastatin decreases coenzyme Q levels in humans. *Proc Natl Acad Sci USA* 1990;87:8931-4.
22. Mortensen SA, Leth A, Agner E, et al. Dose-related decrease of serum coenzyme Q10 during treatment with HMG-CoA reductase inhibitors. *Mol Aspects Med* 1997;18:S137-44.
23. Ghirlanda G, Oradei A, Manto A, et al. Evidence of plasma CoQ10-lowering effect by HMG-CoA reductase inhibitors: A double blind, placebo-controlled study. *J Clin Pharmacol* 1993;33:226-9.
24. Bargossi AM, Grossi G, Fiorella PL, et al. Exogenous CoQ10 supplementation prevents plasma ubiquinone reduction induced by HMG-CoA reductase inhibitors. *Mol Aspects Med*. 1994;15:S187-S193.
25. Food and Drug Administration. Citizen petition to change the labeling for all drugs (Mevacor, Lescol, Pravachol, Zocor, And Advicor) recommending use of 100-200 mg per day of supplemental coenzyme Q10, including cardiomyopathy and congestive heart failure. Available at: <http://www.fda.gov/ohrms/dockets/dailys/02/May02/052902/02p-0244-cp00001-01-vol1.pdf>. Accessed on February 12, 2010.
26. Koumis T, Nathan J, Rosenberg J, Cicero L. Strategies for the prevention and treatment of statin-induced myopathy: Is there a role for ubiquinone supplementation? *Am J Health Sys Pharm*. 2004;61:515-519.
27. Boler JB, Farley TM, Scholler J, Folkers K. Deficiency of coenzyme Q10 in the rabbit. *Int Z Vitaminforsch* 1969;39:281-8.
28. Folkers K, Vadhanavikit S, Mortensen SA. Biochemical rationale and myocardial tissue data on the effective therapy of cardiomyopathy with coenzyme Q10. *Proc Natl Acad Sci USA* 1985;82:901-4.
29. Littarru GP, Tiano L. Bioenergetic and antioxidant properties of coenzyme Q10: recent developments. *Mol Biotechnol*. 2007;37(1):31-7.
30. Tran MT, Mitchell TM, Kennedy DT, et al. Role of Coenzyme Q10 in Chronic Heart Failure, Angina, and Hypertension. *Pharmacotherapy*. 2001; 21:797-806.
31. Morisco C, Trimarco B, Condorelli M. Effect of coenzyme Q10 therapy in patients with congestive heart failure: A long-term, multicenter, randomized study. *Clin Investig*. 1993;71:S134-S146.
32. Baggio E, Gandini R, Plauncher AC, et al. Italian multicenter study on the safety and efficacy of coenzyme Q10 as adjunctive therapy in heart failure. *CoQ10 Drug Surveillance Investigators*. *Mol Aspects Med*. 1994;15:S287-S294.
33. Berman M, Erman A, Ben-Gal T, et al. Coenzyme Q10 in patients with end-stage heart failure awaiting cardiac transplantation: a randomized, placebo-controlled study. *Clin Cardiol*. 2004;27:295-9.
34. Matthews RT, Yang L, Browne S, et al. Coenzyme Q10 administration increases brain mitochondrial concentrations and exerts neuroprotective effects. *Proc Natl Acad Sci USA*. 1998;95:8892-8897
35. Sohal RS, Forster MJ. Coenzyme Q, oxidative stress and aging. *Mitochondrion*. 2007;7 Suppl:S103-11.
36. Gille L, Rosenau T, Kozlov AV, Gregor W. Ubiquinone and tocopherol: dissimilar siblings. *Biochem Pharmacol*. 2008 1;76(3):289-302.

37. Zita C, Overvad K, Mortensen SA, et al. Serum coenzyme Q10 concentrations in healthy men supplemented with 30 mg or 100 mg coenzyme Q10 for two months in a randomised controlled study. *Biofactors*. 2003;18:185-193.
38. James AM, Smith RA, Murphy MP. Antioxidant and prooxidant properties of mitochondrial Coenzyme Q. *Arch Biochem Biophys*. 2004;423:47-56.
39. Turunen M, Olsson J, Dallner G. Metabolism and function of coenzyme Q. *Biochim Biophys Acta*. 2004;1660:171-9.
40. Alleva R, Tomasetti M, Bompadre S, Littarru GP. Oxidation of LDL and their subfractions: kinetic aspects and CoQ10 content. *Mol Aspects Med*. 1997;18 Suppl:S105-12.
41. Langsjoen PH, Langsjoen AM: Overview of the use of CoQ10 in cardiovascular disease. *Biofactors*. 1999;9:273-284.
42. Watts GF, Playford DA, Croft KD, Ward NC, Mori TA, Burke V. Coenzyme Q(10) improves endothelial dysfunction of the brachial artery in Type II diabetes mellitus. *Diabetologia* 2002;45:420-6.
43. Belardinelli R, Mujaj A, Licalaprice F, et al. Coenzyme Q10 and exercise training in chronic heart failure. *Eur Heart J* 2006; 27:2675-81.
44. Groneberg DA, Kindermann B, Althammer M, et al. Coenzyme Q10 affects expression of genes involved in cell signaling, metabolism and transport in human CaCo-2 cells. *Int J Biochem Cell Biol*. 2005;37(6):1208-18.
45. Cooke M, Iosia M, Buford T, et al. Effects of acute and 14-day coenzyme Q10 supplementation on exercise performance in both trained and untrained individuals. *J Int Soc Sports Nutr* 2008; 5: 8 (<http://www.jissn.com/content/5/1/8>)
46. Sato K. Pharmacokinetics of Coenzyme Q10 in Recovery of Acute Sensorineural Hearing Loss due to Hypoxia. *Acta Oto-laryngologica* 1988;105(s458): 95-102.
47. Angeli SI, Liu XZ, Yan D, et al. Coenzyme Q-10 treatment of patients with a 7445A--->G mitochondrial DNA mutation stops the progression of hearing loss. *Acta Otolaryngol*. 2005;125(5):510-2.
48. Khan M, Gross J, Haupt H. A pilot clinical trial of the effects of coenzyme Q10 on chronic tinnitus aurium. *Otolaryngol Head Neck Surg* 2007;136: 72-7.
49. Makhija N, Sendasgupta C, Kiran U, et al. The role of oral coenzyme Q10 in patients undergoing coronary artery bypass graft surgery. *J Cardiothorac Vasc Anesth*. 2008;22(6):832-9.
50. Rosenfeldt F, Marasco S, Lyon W, et al. Coenzyme Q10 therapy before cardiac surgery improves mitochondrial function and in vitro contractility of myocardial tissue. *J Thorac Cardiovasc Surg*. 2005;129(1):25-32.
51. Długosz A, Kuźniar J, Sawicka E, et al. Oxidative stress and coenzyme Q10 supplementation in renal transplant recipients. *Int Urol Nephrol*. 2004;36(2):253-8
52. Rusciani L, Proietti I, Paradisi A, et al. Recombinant interferon alpha-2b and coenzyme Q10 as a postsurgical adjuvant therapy for melanoma: a 3-year trial with recombinant interferon-alpha and 5-year follow-up. *Melanoma Res* 2007; 17(3): 177-83.